



TRANSVERSE MOMENTUM PHYSICS: FROM SPS TO LHC ENERGIES, COMPARISON TO THE DATA ON $\pi^{-12}\text{C}$ -INTERACTIONS AT $P_{\pi^{-}}=40\text{ GeV}/c$

*Y. HUSEYNALIYEV and I. KHAN

Department of Physics, COMSATS Institute of Information Technology, Park Road, Islamabad, Pakistan

(Received June 19, 2009 and accepted in revised form January 05, 2010)

The aim of the presented analysis is to review the transverse momentum physics in SPS CERN, RHIC BNL and LHC CERN energies and investigate inclusive hadron production in $\pi^{-12}\text{C}$ interactions at $P_{\pi^{-}}=40\text{ GeV}/c$. We focus on p_T dependence of the nuclear modification factor (R) for proton and charged π meson production. We demonstrate that the results are in agreement with the SPS and RHIC ones.

Keywords: Nucleus, Transverse momentum, Suppression, Enhancement, $\pi^{-12}\text{C}$, $40\text{ GeV}/c$, Proton, Meson, Heavy ion collision, SPS, RHIC, LHC

1 Introduction

The physics of the transverse momentum (p_T) is of a great interest. J.D Bjorken [1] has shown that the p_T is a constant variable. Then taking into account the Heisenberg's uncertainty principle, $p_T \times A \sim h$, one can find the size of particle emission area (A), where h is the Plank constant. Using p_T one can talk about applicability region of different models (for example, recombination, fragmentation, etc). In a collision the p_T plays crucial role, when:

1. The nuclei barely graze each other ("peripheral" collision), a high- p_T parton (quark and gluon) quickly escapes the medium and it emerges after losing less energy;
2. The collision might be almost head-on ("central" collision), high- p_T partons have larger in-medium path-lengths and thus lose more energy (in average).

Experimental results and theoretical calculations suggest distinction of three p_T regions [2]: bulk (or low), intermediate and high- p_T . It is a rough distinction, because if in 90th years high- p_T was $2\text{ GeV}/c$, with the starting of first heavy-ion collisions (HIC) in Large Hadron Collider (LHC) of European Organization for Nuclear Research (CERN) experiment in 2010 year it is expected to be $13\text{ GeV}/c$. Thus, above mentioned p_T regions will be changed with increasing incident energy. Further we are giving figures which are correct only for today. The bulk region ($p_T < 2\text{ GeV}/c$)

seems to be driven by the thermal properties of the matter created in heavy-ion collisions. Majority of the particles (about 99% of produced hadrons at the Relativistic Heavy Ion Collider (RHIC) of Brookhaven National Laboratory (BNL) have p_T of less than 1 GeV emitted in HIC appear in the low- p_T region. In the intermediate- p_T region ($2 < p_T < 6\text{ GeV}$) one can investigate to the interaction between the hard probes and the bulk matter created in hadron+nucleus (hA), nucleus+nucleus (AA) interactions and HIC. In the high- p_T ($p_T > 6\text{ GeV}/c$) region measured particle spectra are well described by perturbative Quantum Chromo-Dynamics (pQCD) calculations. High- p_T particles suppression is one of the signatures of quark-gluon plasma (QGP), i.e. a thermalized phase in which partons are the relevant degrees of freedom.

The study of hadron production at high transverse momentum (high- p_T) is a sensitive tool to characterize the matter created in ultra-relativistic HIC, and in particular, to detect the possible formation of QGP. Particles at high- p_T result from parton scatterings with high momentum transfer "hard scattering" which can be described by pQCD. The scattered partons will traverse the created medium as they fragment into the observable hadrons. High- p_T particle production in proton+nucleus (pA) and AA interactions was predicted to be suppressed [3] as a consequence of the energy loss of the scattered projectile partons (or hadrons) in the hot and

*Corresponding author : yashar_huseynaliyev@comsats.edu.pk

dense medium (a phenomenon known as “jet quenching”) before the particle production process. Such suppression was observed by experiments at RHIC in central $Au+Au$ and $Cu+Cu$ collisions at a center-of-mass energy of upto $\sqrt{s_{NN}}=200\text{GeV}$ [4]. Particles are suppressed not equally. Another phenomenon, called “Cronin enhancement” [5] is presented in $d+Au$ and peripheral $Au+Au$ collisions. It is believed to be due to multiple nucleons scattering within the nucleus. Neither of these are clearly understood thus require further experimental and theoretical study.

The hard scatterings are still present in the heavy-ion data, but resulting distributions are found to be modified due to medium interactions. Thus, understanding modifications to the high- p_T particle distributions can lead to qualitative conclusions on the energy loss mechanisms within the medium.

The medium effects (suppression and enhancement) which mentioned above, is characterized by the R_{AA} quantity, a term used for the observation of an increased amount of particle production at high- p_T in AA (or pA), compared to the elementary pp reactions [6].

The R_{AA} quantity is called the nuclear modification factor (NMF) and defined by $R_{AA} = \frac{\sigma_{NN}^{inel} d^2N_{AA}/dp_T dy}{\langle N_{coll} \rangle d^2\sigma_{pp}/dp_T dy}$, where $d^2N_{AA}/dp_T dy$ is the invariant yield, $\langle N_{coll} \rangle$ is the mean number of binary (nucleon+nucleon) collisions occurring in central HIC , σ_{NN}^{inel} is the inelastic pp cross section and y is the rapidity, i.e., the relativistic analog of velocity. For perfect binary scaling (in the absence of jet quenching), a value of $R_{AA} = 1$ would be expected. For $R_{AA} < 1$ the spectra is referred to as suppressed, whilst $R_{AA} > 1$ corresponds to an enhancement.

For a more sensitive search of the other nuclear effects the R_{CP} (central to peripheral) modification factor

$$R_{CP} = \frac{d^2N/(2\pi p_T dp_T dy)(central)/N_{coll}(central)}{d^2N/(2\pi p_T dp_T dy)(peripheral)/N_{coll}(peripheral)}$$

is used. If the multiple scattering interpretation of the Cronin enhancement effect is correct, its contribution to the R_{CP} ratio is expected to be

reduced, as it should already be present in the peripheral base line.

It is believed that the emission of hadrons with transverse momentum upto about $6\text{GeV}/c$ in central relativistic HIC is dominated by recombination of partons, where three quarks or a quark/antiquark pair in a densely populated phase space can form a baryon or meson, respectively. High- p_T hadrons originate primarily from the fragmentation of hard scattered partons, in which the single parton with the probability for a parton i hadronize into a hadron h , that carries a fraction $z < 1$ of the momentum of the parent parton. Such hard scatterings occur in the early phase of the reaction, and the transiting partons serve as probes of the strongly interacting matter produced in the collisions. Measurement of identified particles upto the highest possible p_T , establishing the magnitude, p_T and centrality dependence of the suppression pattern, is crucial to constrain the theoretical models and separate contributions of initial and final state effects from the energy loss mechanism [7].

In the presented paper, in conditions of 4π -geometry, we investigate proton and charged π -meson production, the π^-/π^+ ratio in $\pi^{-12}\text{C}$ interactions at $P_{\pi^-}=40\text{GeV}/c$ and compare these to Super Proton Synchrotron (SPS) and RHIC data on proton and heavy ion collision data, seeking similarities in the particle momentum probabilities transverse to the beam axis. A single nuclear modification factor, R is computed for this comparison. The goal is to use the R quantity to learn about the unseen collisions among the quarks and gluons created in this range of collisions.

2. Transverse Momentum Physics in SPS, RHIC and LHC Energies

In this section we have presented the recent development in the transverse momentum physics by measurements of identified hadrons in hadron-nucleus (hA) and nucleus-nucleus (AA) collisions at SPS and $RHIC$ energies and some predictions of LHC experiment.

2.1 Transverse momentum physics in SPS energies

A long-standing puzzle concerns the magnitude of partonic energy loss in nuclear

collisions at the SPS. WA98 data [8] indicated a large enhancement of inclusive π^0 production at $p_T \cdot 3-4\text{GeV}/c$ in central $Pb+Pb$ relative to $p+p$ collisions when normalized per binary collision ($R_{AA}(p_T)$), although a suppression was observed in central relative to peripheral nuclear collisions ($R_{CP}(p_T)$, also normalized per binary collision) by the same experiment.

New measurements of high- p_T hadron suppression at the SPS were presented at the Quark Matter 2005 conference for K_s^0 $R_{CP}(p_T)$ [9] and charged pions and protons [10] compared to radiative energy loss calculations. Introduction of energy loss in a medium, with charged hadron multiplicity scaling as $dN_{ch}/d\eta$ (the density of charged particles per unit of η , where η is the pseudorapidity, a variable which approaches rapidity for particles with mass less than p_T and which is a function only of polar angle), results in good agreement between calculation and data. These new data solve the high- p_T suppression puzzle at the SPS: the medium densities inferred from bulk multiplicity and high- p_T inclusive hadron measurements are consistent. However, significant theoretical uncertainties remain due to a potentially large Cronin effect at these lower energies.

The neutral pion spectra were measured [3] in minimum bias $p+C$ and $p+Pb$ collisions at $\sqrt{s_{NN}}=17.4\text{GeV}$ in the range $0.7 \leq p_T \leq 3.5\text{GeV}/c$. Based on these spectra the NMF for $Pb+Pb$ collisions at CERN SPS energies could be determined using a measured $p+A$ reference. In very central $Pb+Pb$ collisions (0-1%) a significant suppression of high- p_T neutral pions was observed (≈ 0.5) that is reminiscent of the high- p_T hadron suppression observed in $Cu+Cu$ and $Au+Au$ collisions at RHIC. The pion suppression, together with the results at higher energies from RHIC, will allow constraining the energy dependence of hadron suppression as predicted by jet-quenching models.

Transverse momentum spectra upto $4.5\text{GeV}/c$ around mid rapidity of π^\pm , ρ , \bar{p} and K^\pm in $Pb+Pb$ reactions were measured [11] at $\sqrt{s_{NN}}=17.3\text{GeV}$ by the CERN-NA49 experiment. The extracted R_{AA} ratios scaled with the number of binary collisions show a rapid rise with transverse momentum in the covered p_T region. The R_{CP} ratio

stays below one in the π^\pm channel. Interestingly, the R_{CP} ratios stay rather close to the $\sqrt{s_{NN}}=200\text{GeV}$ RHIC results for π^\pm , K^\pm and p in the available p_T region; however the 200GeV π^\pm ratio shows a much larger suppression at $p_T \geq 2\text{GeV}$. The p ratio is almost identical to the 200GeV ratio. The modification factors, when scaled with the number of wounded nucleons, start from one, show a rapid rise with p_T , and the R_{pA} ratios stay below the R_{AA} data over the covered p_T region for both collision energies.

2.2 Transverse momentum physics in RHIC energies

In HIC, high- p_T hadrons, photons and heavy mesons, can be used to probe the soft matter generated in the collision. Initial state production of high- p_T partons is relatively unaffected by the presence of the soft medium, but the partons lose energy when traversing the medium, dominantly due to gluon radiation. The goal of high- p_T measurements is to study these interactions and to use them to measure the density and temperature of the soft matter.

In [12] in $p+p$ collisions, charged pions and (anti-) protons are measured in the range $5.0 \leq p_T \leq 15.0\text{GeV}/c$ and the \bar{p}/π^+ , p/π^- , \bar{p}/p and π^-/π^+ ratios are discussed. The ρ^0 production at high- p_T ($5.0 \leq p_T \leq 10.0\text{GeV}/c$) measured in minimum bias (MB) $p+p$, $Au+Au$ and central $Au+Au$ collisions in the STAR RHIC detector are presented. The ρ^0/π ratio measured in $p+p$ is compared to PYTHIA calculations as a test of pQCD that describes reasonably well particle production from hard processes and also to the ρ^0/π ratio measured in $Au+Au$ collisions. The charged pions, ρ^0 and proton plus anti-proton NMF were also presented. The protons plus anti-protons and the charged pions R_{AA} seem to behave oppositely to what is naively expected from color charge dependence of energy loss. The ρ^0 , charged pions and π^0 NMF are comparable indicating that the fragmentation of vector mesons and pseudo-scalars are similar in $p+p$ and $Au+Au$ collisions.

In paper [13] the π^0 yields for $Cu+Cu$ collisions at $\sqrt{s_{NN}}=22.4$, 62.4 and 200GeV are presented. Reference data for $p+p$ collisions at $\sqrt{s_{NN}}=62.4$

and 200GeV were taken with the same experiment [14]. At $\sqrt{s_{NN}}=22.4\text{GeV}$ a $p+p$ reference was obtained from a parameterization of the world's data on pion production [15]. It is shown that, high- p_T π^0 yields in central $Cu+Cu$ collisions at 62.4 and 200GeV are suppressed, suggesting that parton energy loss is significant, while at 22.4GeV the π^0 yields for $p_T \geq 2\text{GeV}/c$ are not suppressed. The R_{AA} measured in central $Cu+Cu$ at 22.4GeV is consistent with Cronin enhancement alone but does not rule out parton energy-loss effects. These measurements provide a unique constraint for jet-quenching models and demonstrate that parton energy loss starts to prevail over the Cronin enhancement between $\sqrt{s_{NN}}=22.4$ and 62.4GeV.

In paper [16], the J/Ψ spectra at high- p_T ($5 < p_T < 14\text{GeV}/c$) in $p+p$ and $Cu+Cu$ collisions at $\sqrt{s_{NN}}=200\text{GeV}$ are reported. The high- p_T J/Ψ production was found to follow the x_T ($x_T = 2p_T/\sqrt{s}$) scaling with a beam energy dependent factor $\sim (\sqrt{s_{NN}})^{5.6 \pm 0.2}$. Within the errors the power was found to be the same for pion, proton and for J/Ψ , which indicates that the high- p_T J/Ψ production mechanism is closer to parton-parton scattering. The J/Ψ NMF R_{AA} in $Cu+Cu$ increases from low to high- p_T which challenges some models. The average of R_{AA} at $p_T > 5\text{GeV}/c$ is 0.9 ± 0.2 , consistent with no J/Ψ suppression. It implies that high- p_T J/Ψ may be produced from virtual photon or formed outside of the hot interaction region.

The paper [17] exhibits the suppression for direct photons and various mesons (π^0 , η , ϕ , J/Ψ and ω) in central $Au+Au$ collisions at $\sqrt{s_{NN}}=200\text{GeV}$, measured up to $p_T \approx 20\text{GeV}/c$. The comparison of the π^0 suppression in $Au+Au$ and $Cu+Cu$ collisions at $\sqrt{s_{NN}}$ unveils a simple scaling: the suppression only depends on the number of participating nucleons (N_{part}) for the same $\sqrt{s_{NN}}$. Such a scaling with N_{part} is consistent with a parton energy loss picture. More work is needed to understand the different suppression patterns of ϕ and ω mesons as compared to pions and η 's and the scaling of direct photons at high- p_T in central $Cu+Cu$ and $Au+Au$ collisions.

It has also been found at $RHIC$ that at intermediate $p_T \sim 2-6\text{GeV}/c$, the baryon/meson ratio is much larger in HIC than in proton-proton collisions [18]. This could imply that soft production mechanisms, e.g. hadron formation by coalescence of quarks from thermal matter [19], have a significant contribution to baryon (and meson) production up to $p_T \sim 6\text{GeV}/c$.

2.3. Transverse momentum physics in LHC energies

CERN confirmed that the *LHC* will restart in the autumn of 2009. The experimental study of jet quenching has reached a new level of detail and precision, and interpretation of the striking effects that have been observed is currently limited by theoretical uncertainties. *HIC* at the *LHC* will open up a huge new kinematic regime for jet quenching studies, with qualitatively new observables available. While the *LHC* regime may provide better grounds for quantitative theoretical predictions, a complete picture of jet interactions with dense QCD matter must describe all of the phenomena we observe, both at *RHIC* and at the *LHC*. One can only say that with respect to *LHC* energies we have some predictions underlying model calculations (e.g. the *KKT* model of gluon saturation) [20]. In this model, using the calculation and theoretical results, the dependence of the NMF on p_T of pion and open charm and beauty production at *RHIC* and *LHC* are presented. They teach us that at *LHC* at rapidities $y \geq 0$, the NMF is a very slow function of rapidity. This is an anticipated result.

3. Methods and Materials

The NMF, R of the π^\pm -mesons and protons (with momentum $p_p < 1\text{GeV}/c$) as a function of the p_T of these particles in laboratory frame were analyzed by us using the experimental data on π^- ^{12}C -interactions at $P_\pi=40\text{GeV}/c$ (in centre of mass energy it is $\sqrt{s_{NN}}=32\text{GeV}$). The values of R as a ratio of the inclusive spectrum of $1/N_{ev} (dN/dp_T)$, obtained in the events with number of protons $N_p \geq 2$ (we call it multi-nucleon events and number of these events is $N_{ev}=3571$), to the similar ones in the events with $N_p < 2$ ($N_{ev}=5220$) were determined for the first time, i.e.

$$R = \frac{[1/N_{ev} (dN/dp_T)]_{N_p \geq 2}}{[1/N_{ev} (dN/dp_T)]_{N_p < 2}}$$

In conditions of 4π -geometry, the new method is offered to separate the multi-nucleon events for central collisions and to extract more information on the enhancement and suppression phenomenon.

The experimental data were obtained using the 2m propane bubble chamber of the Laboratory of High Energies (LHE) of Joint Institute for Nuclear Research (JINR) (Dubna, Russia) exposed to a beam of relativistic π^- -mesons at the momentum of 40GeV/c at Serpukhov Accelerator. The statistics of the analyzed data consists of 16865 measured $\pi^-+^{12}\text{C}$ inelastic events corresponding to the cross section $179\pm 2\text{mb}$ [21]. The methodological issues connected with the processing of the stereo photos, selection and identification of types of interactions in propane (C_3H_8), reconstruction of the kinematical characteristics of the secondary particles, their identification, and inclusion of corrections due to the loss of particles emitted under large angle to the object plane of the camera are described in details in [22]. The separation of protons and π^+ -mesons was done visually based on their ionization in region $p < 700\text{MeV}/c$ and protons were identified, since a momentum $140\text{MeV}/c$ at which the track starts to be precisely visible (length $> 3\text{cm}$). The analysis on δ -electrons showed that the admixture of protons among the fast positive singly charged particles identified as π^+ -mesons was $12\pm 5\%$ of the measured mean multiplicity of protons [23]. It should be mentioned that π^- -mesons make up the main fraction ($> 95\%$) among the negatively charged particles [23].

4. Results and Discussion

The values of R for the π^\pm -mesons and protons (with momentum $p_p < 1\text{GeV}/c$) as a function of the p_T of these particles in $\pi^-^{12}\text{C}$ -interactions at $P_{\pi^-}=40\text{GeV}/c$ are shown in Figure 1.




Figure 1. $R(p_T)$ dependence for $\pi^-^{12}\text{C}$ -interactions at $P_{\pi^-}=40\text{GeV}/c$. **left** –pions, **right** –proton.

The R values are normalized to unit in our spectra. We can see that for the π^- -mesons and protons, within the error bars $R=1$. But for π^+ -mesons, beginning from $p_T \sim 1.2\text{GeV}/c$, one can observe anomaly behavior with maximum at $p_T \sim 1.2\text{GeV}/c$. This result could be explained with admixture of protons among the fast π^+ -mesons. Another reason could be due to weak criterion in the selection of multi-nucleon events in our experiment. This result needs further analysis.

Figure 2 shows the π^-/π^+ ratio in 0%–12%, MB $\text{Au}+\text{Au}$, and $d+\text{Au}$ collisions [19]. We observe that the π^-/π^+ ratios are consistent with unity in $d+\text{Au}$, MB, and central $\text{Au}+\text{Au}$ collisions. Predictions from a $p\text{QCD}$ based model with and without partonic energy loss are consistent with the data. The behavior of our experimental curve (left graph) is the same that for above mentioned collisions in region $p_T < 3.5\text{GeV}/c$. Some discrepancy maybe is due to weak criterion in the selection of multi-nucleon events in our experiment.




Figure 2. The π^-/π^+ ratios in: (left) $\pi^-^{12}\text{C}$ interactions at 40GeV/c; (right) 12% central (\circ), MB $\text{Au}+\text{Au}$ (\bullet) and $d+\text{Au}$ (\blacktriangle) [24] collisions at $\sqrt{s_{\text{NN}}}=200\text{GeV}$.

The shaded boxes represent the systematic uncertainties in the top 12% central $\text{Au}+\text{Au}$ collisions. The systematic uncertainties for MB $\text{Au}+\text{Au}$ collisions are similar. Curves are the corresponding predictions from a jet quenching model [25].

Figure 3 shows $R(p_T)$ dependence of charged pions ($(\pi^+ + \pi^-)/2$) in different collisions [26]. For comparison our experimental data is given too(left). It is seen that our data approximately consistent with the PHENIX RHIC collisions ones ($\text{Au}+\text{Au}$ at 200GeV/nucleon). One can also see that strong suppression at $\pi^-^{12}\text{C}$ -

interactions are observed, while only small suppression at SPS observed.

Table 1.

No	Experiment	Collision	Energy, $\sqrt{s_{NN}}$, GeV	Particle	Observation	Paper
1	SPS, WA98	Pb+Pb	17.4	π^0	A significant suppression ($.5 \pm .14$) that is reminiscent of the high- p_T suppression observed in Cu+Cu and Au+Au at RHIC.	3
2	SPS, WA98	Pb+Pb	17.3	π^0	large enhancement at $p_T \sim 3-4$ GeV/c in central Pb+Pb relative to p+p collisions; a suppression in central relative to peripheral nuclear collisions	8
3	SPS, NA57	Pb+Pb	17.3	K_s^0 K	R_{CP} is constant at 0.9 for $p_T > 1$ GeV/c and is significantly larger than WA98 for π^0 (≈ 0.6); At top RHIC, the kaon R_{CP} is larger than for π^0 for $p_T \leq 2$ GeV/c, while they are same for higher p_T	9
4	SPS, NA49	Pb+Pb	17.3	p/π K/π Λ/K K/π	enhancement of the baryon/meson ratio for $p_T > 2$ GeV/c; R_{CP} ratio does not show Cronin enhancement for the mesons at larger p_T when using binary collision scaling; a saturation tendency at high p_T is observed.	10
5	SPS, NA49	Pb+Pb	17.3	π^\pm, p, \bar{p} , K^\pm	R_{AA} shows a rapid rise p_T upto 4.5 GeV/c; R_{CP} ratio stays below 1 in the π^\pm channel. R_{CP} ratios stay rather close to the 200 GeV RHIC results, the 200 GeV π^\pm ratio shows a larger suppression at $p_T \geq 2$ GeV; p ratio is almost identical to the 200 GeV ratio.	11
6	RHIC, STAR	Au+Au and p+p	200	π^\pm, ρ^0 , π^0 , \bar{p}	$p+\bar{p}$ & π^\pm R_{AA} seem to behave oppositely to what is naively expected from color charge dependence of energy loss; the ρ^0, π^\pm & π^0 NMF are comparable; fragmentation of mesons are similar, $5 \leq p_T \leq 15$ GeV.	12
7	RHIC, Phenix	Cu+Cu	22.4, 62.4 and 200	π^0	high- p_T yields in central collisions at 62.4 & 200 GeV are suppressed; at 22.4 GeV, yields for $p_T \geq 2$ GeV/c are not suppressed; R_{AA} at 22.4 GeV is consistent with Cronin enhancement; the results demonstrate that suppression starts to prevail over enhancement between 22.4 & 62.4 GeV.	13, 14, 15
8	RHIC, STAR	Cu+Cu and p+p	200	J/ψ	high- p_T production mechanism is closer to parton-parton scattering; R_{AA} in Cu+Cu increases from low to high- p_T ; the average of R_{AA} at $p_T > 5$ GeV/c is 0.9 ± 0.2 , consistent with no J/ψ suppression.	16
9	RHIC, Phenix	Au+Au, Cu+Cu	200	$\pi^0, \gamma, \eta, \phi$, J/ψ	all high- p_T particles are suppressed (except for γ in Cu+Cu); not all mesons are suppressed by the same factor; π^0 suppression only depends on the N_{part} for the same $\sqrt{s_{NN}}$ which is consistent with a parton energy loss picture.	17
10	RHIC, STAR	Au+Au	200	π^\pm, p, \bar{p}	p & \bar{p} less suppressed than π^\pm , in $1.5 \leq p_T \leq 6$ GeV/c; π^-/π^+ & \bar{p}/p ratios show a weak p_T dependence.	18
11	JINR	$\pi^{-12}C$	32	π^\pm, p	for π & p , the $R=1$; for π^+ , the anomaly behavior with maximum at $p_T=1.2$ GeV/c is observed; behavior of π^-/π^+ ratio & $(\pi^+ + \pi^-)/2$ $R(p_T)$ in $p_T < 3.5$ GeV/c is the same that for Au+Au 200 GeV RHIC PHENIX data.	

5. Conclusion and Recommendations

Starting from a discussion on *SPS*, *RHIC* and *LHC*




Figure 3. The $(\pi^+ + \pi^-)/2$ $R(p_T)$ dependence in different collisions.

results, we have tried to motivate the necessity to move from them to $\pi^{-12}C$ -interaction to extract more information on the enhancement and suppression phenomenon. In conditions of 4π - geometry, the new method is offered to separate the multi-nucleon events for central collisions. We demonstrate that the results are in agreement with the *SPS CERN* and *RHIC BNL* ones. The π^-/π^+ ratio of $\pi^{-12}C$ data are consistent with

results obtained for *Au+Au* and *d+Au* collisions. Table 1 presents the papers of the cases concerned, with a description of the experiment, collision, energy, type of particle and observation.

Measurements of identified protons and pions from low- to high transverse momentum have proven to be a valuable tool in understanding the particle production and energy loss mechanisms in relativistic heavy ion collisions. The suppression of hadrons at high transverse momentum leads one to conclude that the partons undergo a large energy loss due to a hot, dense medium created during the collisions. By studying the multi-nucleon events in hadron - nucleus and nucleus - nucleus interactions one can get useful information about collective phenomena. Our results could be used in other hadron - nucleus and nucleus-nucleus interactions with different incident particles and energies, as well as in heavy ion collisions, especially in *SPS*, *RHIC* and *LHC* experiments.

Acknowledgement

It is our pleasure to thank all the members of the international collaboration who took part in data processing from 2 m propane bubble chamber of the Laboratory of High Energies of Joint Institute for Nuclear Research (Dubna, Russia). This work was supported in part by the Higher Education Commission Research Grant N 1-28/HEC/HRD/2007/501.

References

- [1] J.D. Bjorken Fermi Lab- Pub - 82 / 59 - THY (1982).
- [2] arxiv: 0801.2715v1 [nucl-ex] 17 Jan. 2008.
- [3] X-N. Wang and M. Gyulassy, Phys. Rev. Lett. **68** (1992) 1480, arXiv: 0708.2630v2 [nucl-ex] 17 June (2008).
- [4] K. Adcox et al., Nucl. Phys. **A757** (2005) 184, J. Adams et al., Nucl. Phys. **A757** (2005)102, I. Arsene et al., Nucl. Phys. **A757** (2005) 1, B. B. Back et al., Nucl. Phys. **A757** (2005) 28, arXiv: 0606013 [nucl-ex] 16 Jan. 2006.
- [5] J. Adams et al. Phys. Lett. **B637** (2006) 161.

- [6] C. Alt et al., Phys. Rev. **C77** (2008) 034906.
- [7] R. J. Fries et al., Phys. Rev. **C90** (2003) 202303.
- [8] M. Aggarwal et al., Eur. Phys. J. **C23** (2002) 225.
- [9] Quark Matter, 2005 Conference: A. Dainese for NA57 Collaboration, arXiv: nucl-ex/0507012v1 7 July 2005
- [10] Quark Matter, 2005 Conference: A. Laszlo and T. Schuster for NA49 Collaboration, arXiv: nucl-ex/0510054v1 18 Oct. 2005.
- [11] arXiv: 0711.0547v1 [nucl-ex] 4 Nov. (2007).
- [12] arXiv: 0808.3110v1 [nucl-ex] 22 Aug. (2008).
- [13] A. Adare et al., Phys. Rev. Lett., **101** (2008) 162301.
- [14] A. Adare et al. Phys. Rev. **D76** (2007) 051106.
- [15] arXiv:0807.1252v2 [hep-ph] 7 Nov. (2008).
- [16] arXiv: 0804.4846v1 [nucl-ex] 30 April (2008).
- [17] arXiv: 0804.4562v1 [nucl-ex] 29 April (2008).
- [18] B.I. Abelev et al. (STAR), Phys. Rev. Lett. **97** (2006) 152301, nucl-ex/0606003, B. I. Abelev et al. (STAR), Phys. Lett. **B655** (2007) 104, nucl-ex/0703040.
- [19] Z. W. Lin, C. M. Ko, Phys. Rev. Lett. **89** (2002) 202302, nucl-th/0207014 R.C. Hwa, C. B. Yang, Phys. Rev. **C67** (2003) 034902, nucl-th/0211010, R. J. Fries, B. Muller, C. Nonaka, S. A. Bass, Phys. Rev. Lett. **90**, (2003) 202303, nucl-th/0301087.
- [20] arxiv: 0705.2193v2 [hep-ph] 14 Nov. (2007).
- [21] N. Angelov et al., Yad. Fiz. **33** (1981) 1046.
- [22] A. U. Abdurakhimov et al., Yad. Fiz. **18** (1973) 545, A. U. Abdurakhimov et al., Yad. Fiz. **18** (1973) 1251, S. A. Azimov et al., Nucl. Phys. **B107** (1976) 45.
- [23] N. Angelov, K. P. Vishnevskaja and V. G. Grishin et al., JINR Communications, **P1** (1976) 9792.
- [24] J. Adams et al. Phys. Lett. **B616** (2005) 8, J. Adams et al. Phys. Lett. **B637** (2006) 161.
- [25] X. N. Wang, Phys. Rev. **C58** (1998) 2321, Q. Wang et al., Phys. Rev. **C71** (2005) 014903.
- [26] 23rd Winter Workshop on Nuclear Dynamics-Big Sky, February 12-17, 2007: P. Seyboth for NA49 Collaboration.