

DESCRIPTION OF A NEW *CALOGLYPHUS* BERLESE MITE (ACARI: ACARIDAE) INFESTING PULSE IN PAKISTAN

M. Sarwar^{1*} and M. Ashfaq²

¹Pakistan Atomic Energy Commission, Nuclear Institute of Agriculture, Tando Jam-70060,
*Email: dmsarwar64@yahoo.com ²University of Agriculture, Faisalabad-38040, Pakistan.

Abstract

Of the 13 mite species in genus *Caloglyphus* Berlese (family Acaridae) found in Pakistan, *Caloglyphus arbelos* was identified as a new species. The characters of apodeme 3 (*ap3*) which is reduced and broken medially enable the taxonomic separation of this new species from the other described taxa of genus *Caloglyphus* recorded from different geographical areas. A concise key on hypopus traits usable in taxonomy to discern this species from local and global specimens is made available. These studies on accurate diagnosis and identification of pest species are helpful in their successful management.

Keywords: Mite, Taxonomy, *Caloglyphus arbelos*, Pakistan.

1. Introduction

Mites are microscopic organisms belonging to the subclass Acari and class Arachnida in the Phylum Arthropoda. Acaroid mites in the family Acaridae (Tyroglyphidae) including *Caloglyphus* can survive in diverse environments such as storehouses, and human and animal bodies. Their infestation in humans can cause acariosis in several organs including lung, intestine and urinary tract [1]. Infestation of Acaroid mites is a well-known problem for stored grain, often influencing quality and hygienic condition of the grains [2].

A little work has been undertaken on the taxonomy of the *Caloglyphus* mites. Berlese [3] initially proposed and described the genus *Caloglyphus*. Zakhvatkin [4] prepared a comprehensive review of this genus and portrayed 4 new species and redescribed 6 species with improved descriptions. Nesbitt [5, 6], and Samsinak [7] supplemented 1, 3 and 1 new species to this genus, respectively. Mahunka [8, 9, 10, 11] described 2, 1, 2 and 1 new species, respectively. Hughes [12] prepared an excellent compilation of this genus. Tseng and Hsieh [13] redescribed 1 species with improved depiction. Samsinak [14] amended the tribe *Caloglyphini*, re-established the genus *Caloglyphus* and illustrated 1 new species. Channabasavanna et al. [15], Rao et al. [16], and Ashfaq and Chaudhri [17] incorporated 1, 1 and 4 new species, respectively in this genus. Samsinak [18] pointed out 1 new species of the tribe *Caloglyphini*. Zou and Wang [19], Sevastyanov and Radi [20], Sher et al. [21], Klimov [22], and Eraky [23] supplemented 1, 3, 2, 1 and 1 new species, respectively to this genus. Klimov [24] reviewed acarid mites of the tribe *Caloglyphini* with description of a new species. Klimov and Oconnor [25] published phylogeny, historical ecology and systematic of some mites including full descriptions of each taxon, keys and biological information. Oconnor [26] listed two species of genus *Caloglyphus* (= *Sancassania*) phoretic on other species of arthropods. Sarwar and Ashfaq [27, 28], and Sarwar et al. [29, 30] documented 7 new species in genus *Caloglyphus*. Due to the economic importance of *Caloglyphus* mites, the

present project was undertaken to study the systematics and biodiversity of this genus from various climatic regions of Pakistan.

2. Materials and Methods

The present study was carried out in Acarology Research Laboratory, Department of Agricultural Entomology, University of Agriculture, Faisalabad, Pakistan. Samples for mites were collected from Sargodha county (32-05 N, 72-40 E, 218 m) having the mean annual rainfall 360 mm and temperature 24.00 °C. Treating collected samples in the Berlese-Tullgren funnel apparatus, the mites were isolated. An electric light bulb (60 W) was lit, 10 cm above the samples for 24 hours. The collected mites were placed into small glass vials containing 70% alcohol and stored. Mites were slide mounted in Hoyer's medium and covered with cover slip for identification. Using a phase contrast microscope, taxonomical illustrations of specimen were made with a drawing mirror. Measurements of whole body parts of the species are given in micrometer (μ). The terms of body parts and idiosomal chaetotaxy follow Griffiths et al. [31], and terms of leg chaetotaxy and solenidiotaxy follow Griffiths [32].

3. Results and Discussion

Caloglyphus arbelos, new species (Hypopus)

Material examined: Holotype, hypopus, 1 paratype were collected from Sargodha from lentil (*Lens culinaris* M.) on 4.9.1994 (Sarwar) and deposited in Acarology Research Laboratory, Department of Agricultural Entomology, University of Agriculture, Faisalabad.

Description: The description of mite specimen based on the hypopial stage is summarized in ensuing lines: -

Dorsal idiosoma: Body 350 μ long, 290 μ wide, divided into propodosomal and hysterosomal shields. Propodosomal shield 111 μ long, 270 μ wide, dotted antero-medially, 1 pair visible pores, broken striations antero-laterally, remaining shield smooth; setae *vi*, *ve*, *sci*, *sce* and *scs* each 1 pair, simple, 15 μ , 5 μ , 7 μ , 13 μ and 26 μ long, respectively; all setae spaced from each other, *sci-sci* 40 μ , *sce-sce* 88 μ and *sci-sce* 15 μ apart; setae *sci* and *sce* forming a semi-circular line, middle in position. Hysterosomal shield 243 μ long, 290 μ wide, smooth, lateral margins turning towards venter, dotted and striated anteriorly, longitudinal broken striations latero-posteriorly, 11 pairs setae, 9 pairs visible pores. Setae *d1* = *d2* = *d3* = *d4* = 10 μ ; *hi* 19 μ , *he* 12 μ ; *la* 11 μ , *lp1* = *lp2* = 13 μ ; *sae* 55 μ , *sai* 15 μ , long; *d1* - *d1* 84 μ , *d2* - *d2* 59 μ , *d3* - *d3* 74 μ , *d4* - *d4* 84 μ ; *d1* - *d2* 48 μ , *d2* - *d3* 73 μ , *d3* - *d4* 68 μ and *la* - *la* 175 μ apart. Hysterosomal shield anterior margin overlapping propodosomal shield posterior margin upto 33 μ , overlapping area with transverse, broken striations and dots (Fig. 1).

Gnathosoma: Gnathosoma fused pedipalpi, 2 segmented, broad at base, slightly tapering anteriorly, 29 μ long (basal segment 18 μ , distal segment 11 μ , bifurcated anteriorly, 1 pair arista, 43 μ long, 2 pairs small setae (Fig. 3 upper).

Ventral idiosoma: Apodeme 1 (*ap1*) largely Y-shaped, sclerotized, continuing with sternum 1 (*st1*). Sternum 1 (*st1*) 55 μm long, free. Apodeme 2 (*ap2*) free, curved at tip. Apodeme 3 (*ap3*) not meeting apodeme 4 (*ap4*), broken medially. Sternum 2 (*st2*) absent. Apodeme 5 (*ap5*) originating from trochanter IV, making broad, rounded tip anteriorly, not meeting apodeme 4 (*ap4*). Metasternal seta (*mts*) 1 pair, small, each seta inbetween the tips of apodeme 4 (*ap4*) and apodeme 5 (*ap5*) (Fig. 2). Coxal fields I and II, III and IV open, smooth (Fig. 3 middle). Seta *hv* 13 μm long. Genital shield not separated from ventral shield as shown in Fig. 2, smooth, dotted along genital slit. Genital slit elongated, 2 pairs genital suckers and 1 pair paragenital seta (*pr*) mesiad to genital disc (*gdi3*). Genital disc (*gdi3*) oval-shaped without radial striations. Coxal discs *di1* and *di2* present (Fig. 2). Suctorial shield 76 μm long, 97 μm wide, slightly concave antero-medially, broadly rounded posteriorly, dotted, 1 pair anterior suckers, 1 pair anal suckers, both equal in size having dots in between, 1 pair lateral and 1 pair posterior conoids, 2 pairs vestigial suckers towards periphery (Fig. 3 lower), 1 pair suckers below the shield, posteriorly and laterally with sclerotized piece, with pointed ends. Suctorial shield separated from posterior body end by 34 μm , a distance smaller than suctorial shield length (Fig. 2).

Legs: Strong and stout, I-IV measuring 125 μm , 115 μm , 98 μm and 105 μm long, respectively (trochanter base to tarsus tip). Setae and solenidia on legs I-IV segments: coxae 0-0-0-0, trochanters 1-1-1-0, femora 1-1-0-1, genua 3-3-1-0, tibiae 3-3-2-2, tarsi 13-9-8-8. Tarsi I and II 41 μm and 38 μm long, respectively. Seta *vF* on femora I, II and IV 42 μm , 40 μm and 29 μm long, respectively, absent on femur III. Seta *e* on tarsi I-IV 35 μm , 26 μm , 28 μm and 30 μm in length, respectively. Seta *mG* on genua I and II each lancet-like; *hT* on tibiae I and II each lancet-like, 15 μm , 18 μm , 20 μm and 20 μm long, respectively. Seta \acute{o} on genua I and II, a simple seta 36 μm and a solenidion 11 μm long, respectively. Tarsi I and II each with a solenidion *w1* 25 μm long. Tarsi III and IV short and stout. Seta \ddot{o} on tibiae I and II 70 μm and 53 μm long, respectively. Seta *ba* on tarsus I, 22 μm long. Tarsi I-IV provided with 1 spoon-shaped + 5 leaf-like; 1 spoon-shaped + 4 leaf-like; 1 lancet-like + 3 leaf-like; 1 spoon-shaped + 3 leaf-like setae, respectively (Fig. 2).

Remarks: Much morphological variations have been found in *Caloglyphus arbelos* and other species of genus *Caloglyphus* recorded so far in the world. The new species seems close to *Caloglyphus agrios* Sarwar et al. [29] with the following differing characters: -

1. Propodosomal shield without visible pores and hysterosomal shield with 3 pairs visible pores in *C. agrios* but propodosomal shield with 1 pair and hysterosomal shield with 9 pairs visible pores in this species.
2. Sternum 2 (*st2*) presents in *C. agrios* but absent in this species.
3. Ventral shield separated from genital shield in *C. agrios* but not separated in this species.
4. Coxal discs (*di1*, *di2*) conoids in *C. agrios* but not so in this species.
5. Genital disc (*gdi3*) kidney-shaped in *C. agrios* but oval-shaped in this species.
6. Coxal field III closed in *C. agrios* but open in this species.

Key to Species of Genus *Caloglyphus* (Hypopus)

1 Sternum 2 (<i>st2</i>) present.....	2
- Sternum 2 (<i>st2</i>) absent.....	26
2 Sternum 2 (<i>st2</i>) meeting genital shield posteriorly.....	3

- Sternum 2 (<i>st2</i>) not meeting genital shield posteriorly.....	19
3 Apodeme 2 (<i>ap2</i>) meeting apodeme 3 (<i>ap3</i>).....	4
- Apodeme 2 (<i>ap2</i>) not meeting apodeme 3 (<i>ap3</i>).....	7
4 Dorsum dotted; sternum 1 (<i>st1</i>) bifid posteriorly.....	<i>C. captus</i> Sarwar and Ashfaq
- Dorsum not dotted; sternum 1 (<i>st1</i>) not bifid posteriorly.....	5
5 Propodosomal setae (<i>sci, sce</i>) forming a straight line.....	<i>C. austerus</i> Sarwar et al.
- Propodosomal setae (<i>sci, sce</i>) not forming a straight line.....	6
6 Apodeme 2 (<i>ap2</i>) meeting apodeme 4 (<i>ap4</i>); metasternal seta (<i>mts</i>) present.....	<i>C. spinipes</i> Mahunka
- Apodeme 2 (<i>ap2</i>) not meeting apodeme 4 (<i>ap4</i>); metasternal seta (<i>mts</i>) absent.....	<i>C. rodionovi</i> A. Z.
7 Apodeme 3 (<i>ap3</i>) meeting apodeme 4 (<i>ap4</i>).....	12
- Apodeme 3 (<i>ap3</i>) not meeting apodeme 4 (<i>ap4</i>).....	8
8 Gnathosomal lateral margins parallel.....	9
- Gnathosomal lateral margins not parallel.....	11
9 Sternum 1 (<i>st1</i>) bifid posteriorly; paragenital seta (<i>pr</i>) bifid.....	<i>C. multaniensis</i> Ashfaq and Chaudhri
- Sternum 1 (<i>st1</i>) not bifid posteriorly; paragenital seta (<i>pr</i>) not bifid.....	10
10 Tarsus I with 4 leaf-like setae; external ventral seta of genu II simple.....	<i>C. tshernyshevi</i> Zakhvatkin
- Tarsus I with 5 leaf-like setae; external ventral seta of genu II spine-like.....	<i>C. agrios</i> Sarwar et al.
11 Setae <i>sci</i> and <i>sce</i> forming straight line; apodemes 4 (<i>ap4</i>) not meeting medially.....	<i>C. opacatus</i> Ashfaq and Chaudhri
- Setae <i>sci</i> and <i>sce</i> not forming straight line; apodemes 4 (<i>ap4</i>) meeting medially.....	<i>C. trigonellum</i> Sher et al.
12 Gnathosoma notched posteriorly.....	13
- Gnathosoma not notched posteriorly.....	14
13 Setae <i>sci</i> and <i>sce</i> of equal size; apodemes 4 (<i>ap4</i>) meeting medially.....	<i>C. merisma</i> Ashfaq and Chaudhri
- Setae <i>sci</i> and <i>sce</i> not of equal size; apodemes 4 (<i>ap4</i>) not meeting medially.....	<i>C. hadros</i> Sarwar et al.
14 Dorsal body setae simple.....	15
- Dorsal body setae serrated.....	<i>C. oviformis</i> Mahunka
15 Gnathosoma much below the anterior margin of propodosoma; coxal field III open.....	<i>C. moniezi</i> A.Z.
- Gnathosoma not much below the anterior margin of propodosoma; coxal field III not open...	16
16 Gnathosoma parallel laterally; distal fork separated from basal joint.....	18
- Gnathosoma not parallel laterally; distal fork not separated from basal joint.....	17
17 Hysterosomal shield smooth; trochanter I without seta; ϕ on genu II spine-like.....	<i>C. muscarius</i> Sevastyanov and Radi
- Hysterosomal shield dotted; trochanter I with 1 seta; ϕ on genu II a solenidion.....	<i>C. kenos</i> Sarwar and Ashfaq.
18 Hysterosomal shield smooth; sternum 1 (<i>st1</i>) not bifid posteriorly; coxal discs (<i>di1, di2</i>) conoids.....	<i>C. bradys</i> Sarwar et al.

- Hysterosomal shield dotted; sternum 1 (<i>st1</i>) bifid posteriorly; coxal discs (<i>di1</i> , <i>di2</i>) not conoids.....	<i>C. faisalabadiensis</i> Sher, Asfaq and Parvez
19 Ventral shield separated from genital shield.....	22
- Ventral shield not separated from genital shield.....	20
20 Apodemes 4 (<i>ap4</i>) meeting medially.....	<i>C. polyphyllae</i> Zakhvatkin
- Apodemes 4 (<i>ap4</i>) not meeting medially.....	21
21 Suctorial shield rounded.....	<i>C. mandzhur</i> Zakhvatkin
- Suctorial shield longer than wide.....	<i>C. rhizoglyphoides</i> Zakhvatkin
22 Gnathosoma with 1 pair small setae.....	23
- Gnathosoma not with 1 pair small setae.....	25
23 Coxal disc (<i>di1</i>) replaced by a seta.....	<i>C. mycophagus</i> Megnin
- Coxal disc (<i>di1</i>) not replaced by a seta.....	24
24 Sternum 2 (<i>st2</i>) free from either sides; coxal field III open.....	<i>C. baloghi</i> Mahunka
- Sternum 2 (<i>st2</i>) free posteriorly; coxal field III closed.....	<i>C. conus</i> Mahunka
25 Apodeme 2 (<i>ap2</i>) meeting apodeme 3 (<i>ap3</i>); suctorial shield with radial striations posteriorly.....	<i>C. bifurcatus</i> Mahunka
- Apodeme 2 (<i>ap2</i>) not meeting apodeme 3 (<i>ap3</i>); suctorial shield without radial striations posteriorly.....	<i>C. forficularis</i> Sevastyanov and Radi
26 Gnathosoma extended beyond the body.....	27
- Gnathosoma not extended beyond the body.....	29
27 Apodemes 4 (<i>ap4</i>) meeting medially.....	<i>C. morosus</i> Ashfaq and Chaudhri
- Apodemes 4 (<i>ap4</i>) not meeting medially.....	28
28 Coxal field III open; metasternal seta (<i>mts</i>), paragenital seta (<i>pr</i>) and seta <i>hv</i> present.....	<i>C. arbelos</i> , n. sp.
- Coxal field III closed; metasternal seta (<i>mts</i>), paragenital seta (<i>pr</i>) and seta <i>hv</i> not present.....	<i>C. berleseii</i> Michael
29 Apodeme 3 (<i>ap3</i>) meeting apodeme 4 (<i>ap4</i>).....	32
- Apodeme 3 (<i>ap3</i>) not meeting apodeme 4 (<i>ap4</i>).....	30
30 Gnathosoma reduced; coxal field III opened.....	<i>C. caroli</i> Channabasavanna <i>et al.</i>
- Gnathosoma well developed; coxal field III closed.....	31
31 Coxal field II open; legs strong and stout.....	<i>C. geotruperum</i> Zakhvatkin
- Coxal field II closed; legs fat and small.....	<i>C. spinatarsus</i> Hermann
32 Coxal field III open; genital disc (<i>gdi3</i>) and suctorial shield with radial striation.....	<i>C. clemens</i> Sarwar and Ashfaq
- Coxal field III closed; genital disc (<i>gdi3</i>) and suctorial shield without radial striation.....	<i>C. cingentis</i> Sarwar and Ashfaq

4. Conclusions

Our results have helped to clarify the taxonomic status of one new *Caloglyphus* mite detected infesting the lentil. It seems distinct from already existing taxa of the genus. The characters of apodeme 3 (*ap3*) which is reduced and broken medially enable the taxonomic separation of this new species from the other described taxa of genus *Caloglyphus* recorded in different geographical areas of Pakistan. These results would be helpful in mite control efforts because any successful management of a pest starts with its correct diagnosis and identification.

Acknowledgments

We thank the numerous friends and colleagues who helped us in mite collections.

References

- [1] Boquete, M., Carballada, F., Armisen, M., Nieto, A., Martin, S., Polo, F. and Carreira, J. 2000. Factors influencing the clinical picture and the differential sensitization to house dust mites and storage mites. *J. Investig. Allergol. Clin. Immunol*, 10, 229-234.
- [2] Li, C.P., Cui, Y.B., Wang, J., Yang, Q.G. Tian and Y. 2003. Acaroid mite, intestinal and urinary acariasis. *World Journal of Gastroenterology*, 9 (4), 874-877.
- [3] Berlese, A. 1923. Centuria sesta di Acari Nuovi. *Redia*, 15, 237-262.
- [4] Zakhvatkin, A. A. 1941. *Fauna of USSR Arachnoidea VI (1) Tyroglyphoidea (Acari)*. Zoological Institute of the USSR Academy of Sciences, New Series, No. 28. English Translation 1959, Rateliffe, A., and Hughes, A. M., American Institute of Biological Sciences, 573 pp.
- [5] Nesbitt, H. H. J. 1944. Three new mites of the subfamily Rhizoglyphinae. *Canadian Entomologist*, 76 (2), 21-27.
- [6] Nesbitt, H. H. J. 1949. Six new Mexican mites of the sub family Rhizoglyphinae, Acarina. *Pan Pacific Entomologist*, 25 (2), 57-70.
- [7] Samsinak, K. 1966. Die Neuerrichtung der Gattung *Cosmoglyphus* Oudmans 1932 gleichzeitig ein Beitrag zum Problem der "Copra itch". *Zoologischer Anzeiger*, 176 (1), 27-42.
- [8] Mahunka, S. 1973. Auf insekten lebende Milben (Acari: Acarida, Tarsonemida) aus Afrika II. *Acta Zoologica Hungrica*, 19 (3-4), 289-337.
- [9] Mahunka, S. 1974. Auf insekten lebende Milben (Acari: Acarida, Tarsonemida) aus Afrika III. *Acta Zoologica Hungrica*, 20 (1-2), 137-154.
- [10] Mahunka, S. 1978. Schizoglyphidae fam. n. and new taxa of Acaridae and Anoeidae (Acari: Acarida). *Acta Zoologica Hungrica*, 24 (1-2), 107-131.
- [11] Mahunka, S. 1979. The examination of myrmecophilous Acaroidea mites based on the investigations of Dr. C. W. Rettenmeyer (Acari: Acaroidea) II. *Acta Zoologica Hungrica*, 25, 311-356.
- [12] Hughes, A. M. 1976. *The Mites of Stored Food and Houses. Technical Bulletin, No. 9*, Ministry of Agriculture Food and Fisheries, London, 400 pp.
- [13] Tseng Y. H. and Hsieh, S. A. 1976. A new record of acarid mite *Caloglyphus mycophagus* (Megnin) from Taiwan (Acarina: Astigmata). *Taiwan Sugar Research Institute*, No. 74, 47-52.
- [14] Samsinak, K. 1980. *Caloglyphus rodriguezi* new species with taxonomic remarks on the tribe *Caloglyphini* (Acari: Acaridae). *Mitteilungen aus dem Zoologischen Museum Berlin*, 56 (2), 201-206.
- [15] Channabasavanna, G. P., Krishna, N. S. Rao and Ranganath, H. R. 1981. A new *Caloglyphus* (Astigmata: Acaridae) from poultry litter in India with taxonomic comments on the genus. *Indian Journal of Acarology*, 6 (1/2), 57-63.
- [16] Rao, N. S. K., Ranganath, H. R., Channabasavanna, G. P., Krishna, N. S. Rao and Rao, N. S. K. 1982. *Caloglyphus karnatakaensis* sp. Nov. (Acari: Acaridae) from India with

- taxonomic comments on the genus *Caloglyphus*. *Indian Journal of Acarology*, 7 (1), 37-43.
- [17] Ashfaq, M. and Chaudhri, W. M. 1983. Four new (Hypopi) species of the genus *Caloglyphus* Berlese from Pakistan (Acarina: Acaridae). *Pakistan Entomologist*, 5 (1-2), 61-78.
- [18] Samsinak, K. 1988. *Sancassania ultima* a new mite of the tribe *Caloglyphini* (Acari: Acaridae). *Entomologie Mitteilungen aus dem Zoologischen Museum Hamburg*, 9 (133), 159-164.
- [19] Zou, P. and Wang, X. Z. 1989. A new species and two new records of Acaridae associated with edible fungi from China (Acarina: Acaroidea). *Acta Agriculturae, Shanghai*, 5 (3), 21-24.
- [20] Sevastyanov, V. D. and Radi, G. K. K. K. 1991. New species of the mite family Acaridae (Sarcoptiformes) from Lower Egypt. *Entomology Review*, 8, 139-146.
- [21] Sher, F., Ashfaq, M. and Parvez, A. 1991. Two new (hypopi) species of genus *Caloglyphus* Berlese (Acarina: Acaridae) from Pakistan. *Pakistan Entomologist*, 13 (1-2), 27-34.
- [22] Klimov, P. B. 2000. A review of acarid mites of the tribe *Caloglyphini* (Acaridae: Acariformes) with description of a new genus and species from Siberia and Russian Far East. *Vestnik Zoologii Ukraine*, 34 (4-5), 27-35.
- [23] Eraky, S. A. 1999. Five new hypopial nymphs (Acari: Acaridae and Histiostomatidae) described from different habitats. *Folia Entomologica Hungarica*, 60, 45-56.
- [24] Klimov, P. B. and Oconnor, B. M. 2003. Phylogeny historical ecology and systematics of some mushroom associated mites of the genus *Sancassania* (Acari: Acaridae) with new generic synonymies. *Invertebrate Systematics*, 17, 469-514.
- [25] Klimov, P. V. 1996. A new species of acarid mite from the genus *Caloglyphus* (Acari: Acaridae) from the Russian Far East. *Zoologicheskii Zhurnal*, 75 (4), 613-619.
- [26] Oconnor, B. M. 2003. Systematics and ecology of North American bee-associated mites: potential threats to native and introduced pollinators. *Ann Arbor Michigan, USA*, 1079-1098.
- [27] Sarwar, M. and Ashfaq, M. 2004. Two new *Caloglyphus* Berlese mites (Astigmata: Acaridae) recorded in Pakistan. *Pakistan Journal of Scientific and Industrial Research*, 47 (6), 455-461.
- [28] Sarwar, M. and M. Ashfaq. 2006. A new mite pest (Acarina: Acaridae) detected from stored commodity in the Punjab province of Pakistan. *Acarologia*, XLVI, 1-2, 115-120.
- [29] Sarwar, M., Ashfaq, M. and Akbar, S. 2005. Numerical taxonomy of two new mites species of the genus *Caloglyphus* Berlese (Acaridae) from Pakistan. *Pakistan Journal of Scientific and Industrial Research*, 48 (5), 345-353.
- [30] Sarwar, M., M. Ashfaq and S. Nadeem. 2009. On the identity of new acarid mites in genus *Caloglyphus* Berlese occurring in Asian expanse (Pakistan) (Acarina: Acaridae). *Journal of Agriculture and Biological Sciences*, 1 (1), 38-47.
- [31] Griffiths, D. A., Atyeo, W. T., Norton, R. A. and Lynch, C. A. 1990. The idiosomal chaetotaxy of astigmatid mites. *Journal of Zoology*, 220, 1-32.
- [32] Griffiths, D. A. 1970. A further systematic study of the genus *Acarus* L. 1758 (Acaridae: Acarina) with a key to species. *Bulletin of British Museum (Natural History), Zoology Series*, 19, 85-118.




Fig. 1 showing Dorsal view of *C. arbelos* n. sp.





Fig. 2 showing Ventral view of *C. arbelos* n. sp.



Figs. 3, Gnathosoma (upper), Coxal apodemes (middle), Suctorial shield (lower) of *C. arbelos* n. sp.